

## Clinical Guide to Surgical Prioritisation During the Coronavirus Pandemic

This Guide was produced at the request of NHS England at the start of the pandemic. It is written by specialists in the procedures listed and is updated at least every month. It sets out what clinicians view as the relative priorities of conditions at the time it is posted.

It is essential that all patients listed in any category are regularly, clinically reviewed to ensure their condition is not changing and in need of re-prioritising.

The relative priorities between cases listed in the same time frame will need to be decided locally in relation to facilities available and local Covid conditions.

The 'RPM' form, included in the footer of the guide, is designed to help review and reprioritise cases in p1b-4.

The Guide is a short term expedient to the pandemic and not for long term use.

Both the Guide and the RPM are available to down load at (https://fssa.org.uk/covid-19\_documents.aspx)

With thanks to all the Surgical Associations, which have contributed and to the RCOG, RCOphth, RCPSG, RCSEd, RCSEng and RCSI

The Association of Surgeons of Great Britain & Ireland

The British Association of Oral & Maxillofacial Surgeons

ENT -UK

The British Association of Plastic, Reconstructive & Aesthetic Surgeons
The Society for Cardiothoracic Surgery in Great Britain & Ireland

The British Association of Paediatric Surgeons

The British Association of Urological Surgeons

The British Orthopaedic Association

The Society of British Neurological Surgeons

The Vascular Society of Great Britain & Ireland

		e performed in <24 ho		- see notes below									
General surgery (including	Emergency laparotomy -	Appendicectomy - complicated/	Intra-abdominal trauma - unsuitable	Drainage of localised sepsis/necrosis - not	oesophagus/	Acute airway obstruction - thyroid	All solid organ transplants						
oesophago-gastric, HPB, coloproctology,	Peritonitis	unresponsive to conservative Rx	for/not responding to conservative Rx	responding to conservative Rx (antibiotics/	stomach - with survivable mediastinitis/								
breast, endocrine)	Perforation Ischaemia			Interventional radiology)	peritonitis								
	Necrotising fasciitis												
	Small and large bowel obstruction with concerning												
	features of incipient ischaemia/												
	perforation  Post-operative												
	complications (e.g. anastomotic leaks)												
	Bleeding - not suitable for/ responding to												
	endoscopic/control/ interventional radiology												
OMFS		Dental Sepsis - not	Orbital Compartment	Jaw Dislocation - not responding to									
	trauma not responsive to conservative Rx	conservative Rx and		conservative Rx									
	(reduction + IR)	Signit brain.	to signi										
Reconstructive plastic surgery	Major burns - Airway	Chemical burns - especially Eye/	Necrotising Fasciitis - any site	Soft tissue infection - any site (especially	re-implantation/	Washout open wound/fractures/	Removal of prosthesis/expander						
including burns and hands	management/ resuscitation/ escharotomies/	Hydrofluoric acid >2%		closed compartments/ joints/prostheses)	failing free flap - any site	infected/grossly contaminated (human/animal/	for fulminant infection						
	amputations/Toxic Shock			not responding to conservative Rx		contaminated) wounds - any site							
Urology	Renal obstruction with infection - not	Renal/ureteric trauma requiring	Bladder trauma requiring open	Genital trauma/ testicular torsion/	Fournier's gangrene	Haematuria/ uncontrolled	Insertion of catheter under GA						
	responding to conservative Rx	open surgery	surgery	amputation/priapism (>24hrs)		haemorrhage - causing	under div						
						haemodynamic instability and unresponsive to							
Trauma and	Fractures -	Infection -	Dislocated joints	Compartment	Spinal Trauma with	conservative Rx  Acute spinal cord	Cauda Equina						
orthopaedics (including spinal surgery)	Open	Septic arthritis - (natural or prosthetic		syndrome	instability and or neurological dysfunction	compression - with neurological dysfunction -	Syndrome - Clinically and radiologically						
	Neurovascular +/- Skin compromise	joint) Other metalwork				including MSCC	confirmed.						
	Hip/femoral shaft  Long bone/Pelvic +/-	(including spine)											
	Spinal fixation in polytrauma												
ENT	Airway obstruction -		Nasal/ear button	Life threatening	Orbital cellulitis	Uncontrolled	Sinus surgery for						
	Cancer/Foreign body/Sepsis	vascular/visceral/ airway injury	battery removal	middle ear conditions		epistaxis	impending catastrophe/failure to respond to						
Neurosurgery (including spinal	Traumatic Brain injury - unsuitable	Intra-cranial haemorrhage - not	Acute raised Intra cranial pressure/	Cauda Equina Syndrome -	Myelomeningocoele		medical Rx						
surgery)	for conservative RX	responding to conservative RX	Hydrocephalus	Clinically and radiologically confirmed.									
			suitable for conservative Rx	SSIIIGU.									
Cardiothoracic	Ruptured bronchus	Myocardial infarction -	Empyema with	Aortic dissection	Acute presentation	Acute mitral valve	Chest Trauma						
surgery		imminent death	sepsis		of ventricular septal defect	disease							
Vascular surgery	Vascular injury/ occlusion - Limb/	Uncontrolled external	Ruptured AAA	Diabetic foot sepsis									
	Gut/SVC	haemorrhage - any site/source											
Paediatric general	Neonatal	Emergency	Emergency	Appendicectomy -	Thoracotomy/Chest	Strangulated	Acute Scrotal	Trauma	Trauma Laparotomy	Removal of Infected	Renal Obstruction -	Bladder outlet or	Urosepsis - not
and urological surgery (see also urology)	Malformations -  Oesophageal	Laparotomy - (Neonatal) -	laparotomy - (Infant/child)		Drain Insertion/Video Assisted Thorascopic Surgery	inguinal hernia	Exploration (suspected Testicular Torsion)	Thoracotomy		Central Line	i) Infection/pain - not responding to	urethral obstruction	responding to conservative Rx.
	Atresia,  Gastroschisis,	Necrotising Entero- Colitis (NEC),	Peritonitis Perforation	Conscivative 11X	(VATS) for Empyema		Tostional Torsion)				conservative Rx  ii) Impaired renal		
	Anorectal	Perforation,	Intussusception								function		
			"" taccaccop "" o"										
	Malformations	Malrotation	Ischaemia								iii) Single kidney		
	Malformations	Malrotation											
	Malformations	Malrotation	Ischaemia  Necrotising fasciitis										
	Malformations	Malrotation	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g.										
	Malformations	Malrotation	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative										
Paediatric orthopaedic	Malformations  Septic arthritis/ osteomyelitis	Fractures -	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/	Compartment syndrome									
	Septic arthritis/	Fractures - Open Neurovascular	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)										
orthopaedic surgery (including spinal surgery)	Septic arthritis/ osteomyelitis	Fractures - Open Neurovascular compromise +/-Skin compromise	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints	syndrome	Namata	Infant Dight heart	Infant Doguveitant	Child Left beaut	Child Deguratent				
orthopaedic surgery (including	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions -	Fractures - Open Neurovascular compromise +/-Skin compromise	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -	syndrome  Neonate - Shunt/ stent -	Neonate - Arrhythmia	Infant - Right heart obstructive lesions -	Infant - Regurgitant lesions -	Child - Left heart obstructive lesions -	Child - Regurgitant lesions -				
orthopaedic surgery (including spinal surgery)	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions -  HLHS (restrictive/ intact atrial septum)	Fractures -  Open  Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions -  PA-IVS	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/		obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to							
orthopaedic surgery (including spinal surgery)	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions -  HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS	Neonate - Shunt/ stent -  Profound hypoxaemia/	Arrhythmia  CHB not responding	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
orthopaedic surgery (including spinal surgery)	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions -  HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/	Arrhythmia  CHB not responding	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
orthopaedic surgery (including spinal surgery)	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions -  HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/	Arrhythmia  CHB not responding	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
orthopaedic surgery (including spinal surgery)	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions -  HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD  Tetralogy of Fallot Critical pulmonary stenosis	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/	Arrhythmia  CHB not responding	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
orthopaedic surgery (including spinal surgery)  Paediatric cardiac surgery  Obstetrics and Gynaecology	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions -  HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to	syndrome  Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)	Arrhythmia  CHB not responding	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
orthopaedic surgery (including spinal surgery)  Paediatric cardiac surgery  Obstetrics and Gynaecology (including urogynaecology, pregnancy,	Septic arthritis/osteomyelitis  Neonate - Left heart obstructive lesions -  HLHS (restrictive/intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/Laparoscopy  Miscarriage with bleeding requiring	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
orthopaedic surgery (including spinal surgery)  Paediatric cardiac surgery  Obstetrics and Gynaecology (including urogynaecology,	Septic arthritis/osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation  b) Females - before	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen -	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions -  HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis,	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation  b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe pre-eclampsia, etc.)	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions -  HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/ 9+6	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation  b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding	Fractures - Open  Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS  PA-VSD  Tetralogy of Fallot  Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery  Emergency Caesarean  Instrumental delivery  Perineal repair  Manual removal of placenta  Cervical cerclage  Emergency laparotomy/	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6)	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation  b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma	Fractures - Open  Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS  PA-VSD  Tetralogy of Fallot  Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery  Emergency Caesarean  Instrumental delivery  Perineal repair  Manual removal of placenta  Cervical cerclage  Emergency laparotomy/	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/9+6 weeks [International for medical abortion at home)/12-14 weeks	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation  b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelve-vaginal haematoma)  Ectopic pregnancy	Fractures - Open  Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS  PA-VSD  Tetralogy of Fallot  Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery  Emergency Caesarean  Instrumental delivery  Perineal repair  Manual removal of placenta  Cervical cerclage  Emergency laparotomy/	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe pre-eclampsia, etc.)  Approaching legal threshold (23+6 weeks fe Enll and & Wales]/114-6 weeks [Scotland] for medical abortion at	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation  b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP	Fractures - Open  Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS  PA-VSD  Tetralogy of Fallot  Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery  Emergency Caesarean  Instrumental delivery  Perineal repair  Manual removal of placenta  Cervical cerclage  Emergency laparotomy/	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home/ 12-14 weeks [Scotland] for medical for medical abortion at home/ 12-14 weeks where in the provided by local not provided by local not provided by local not preparation has	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation  b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy - (heavy bleeding requiring evacuation)  requiring evacuation,	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks [England & Wales]/11+6 weeks	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy - (heavy bleeding	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/ 9+6 weeks [England & Wales]/11+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical for at home/ 12-14 weeks (Scotland] for medical for at home/ 12-14 weeks (Scotland)	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
orthopaedic surgery (including spinal surgery)  Paediatric cardiac surgery  Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive	Septic arthritis/osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy - (heavy bleeding requiring evacuation)  requiring evacuation,	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe pre-eclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/ 9+6 weeks [England & Wales]/11+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home/ 12-14 weeks where in a more in	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy Complications of TOP  Molar pregnancy - (heavy bleeding requiring evacuation hysterectomy)	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/ 9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home/ 12-14 weeks (Scotland] for medical abortion at home/ 12-14 weeks (Scotland) for medical abortion at home/ 12-14 weeks	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy - (heavy bleeding requiring evacuation, hysterectomy)  Acute risk to sight/ life (e.g. penetrating injuries/globe rupture/orbital haemorrhage/burns	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  Oncology - Ruthenium plaque	Ischaemia Necrotising fasciitis Bleeding (not responding to conservative Rx) Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions - TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care - Miscarriage - bleeding and unstable Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/ 9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home)/ 12-14 weeks where procedure and proce	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy (reavy bleeding requiring evacuation, hysterectomy)  Acute risk to sight/ life (e.g. penetrating injuries/globe rupture/orbital	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  Oncology - Ruthenium plaque	Ischaemia Necrotising fasciitis Bleeding (not responding to conservative Rx) Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions - TGA (hypoxaemia for BAS/Intact IVS for ASO) TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care - Miscarriage - bleeding and unstable Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/ 9+6 weeks [Ingland & Wales]/11+6 weeks [Scotland] for medical abortion at home)/ 12-14 weeks where procedy ure not provided by local NHS beyond this)  Cases where cervical preparation has been administered (misoprostol/osmotic dilators/mifepristone) Feticide (approaching legal limit)  Vitreous biopsy/ antibiotic injection - suspected	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Arrhythmia  CHB not responding to medical Rx.  Paediatric/ adolescent  Imperforate hymen - incision and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy Complications of TOP  Molar pregnancy (heavy bleeding requiring evacuation, hysterectomy)  Acute risk to sight/ life (e.g. penetrating injuries/globe rupture/orbital haemorrhage/burns infection/fractures/ lid lacerations)	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  Oncology - Ruthenium plaque removal.	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home)/ 12-14 weeks where procedure not provided by local NHS beyond this)  Cases where cervical provided by local NHS beyond this)  Cases where cervical provided by local NHS beyond this)  Cases where cervical provided by local NHS beyond this)  Cases where cervical provided by local NHS beyond this)  Visteoretinal  Vitreous biopsy/ antibiotic injection - suspected endophthalmitis	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine - a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	Paediatric/adolescent  Imperforate hymen - incision and drainage	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)  PLEASE NOTE: More detailed	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy (heavy bleeding requiring evacuation, hysterectomy)  Acute risk to sight/ life (e.g. penetrating injuries/globe rupture/orbital haemorrhage/burns infection/fractures/	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  Oncology - Ruthenium plaque	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/ bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/ 9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home)/ 12-14 weeks where procedure not provided by local NHS beyond this)  Cases where cervical preparation has been apministered (misoproshifosmotic dilators/mifepristone)  Feticide (epiparation legal limit)  Vitreoretinal  Vitreous biopsy/ antibiotic injection - suspected endophthalmitis	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine -  a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	e) Safeguarding issues must be	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)  PLEASE NOTE: More detailed specialty specific guidance can be found on the NHSE	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy Complications of TOP  Molar pregnancy - (heavy bleeding requiring evacuation, hysterectomy)  Acute risk to sight/ life (e.g. penetrating inuptive/glober haemorinhgractures/ lid lacerations)  a) THIS DOCUMENT WILL BE REVIEWED MONTHLY	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  b) This Prioritisation is about 'WHEN and not BY Whom'. Space does not	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/ bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home/r 12-14 were not provided by local NHS beyond this)  Cases where cervical preparation has been administered (misoprostol/osmotic dilators/mifepristone)  Feticide (approaching legal limit)  Vitreoretinal	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine - a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	e) Safeguarding issues must be considered in all those attending with trauma and	obstructive lesions -  Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)  PLEASE NOTE: More detailed specialty specific guidance can be found on the NHSE website https://www.england.nhs.uk/coronavirus/	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (requiring evacuations hysterectomy)  Acute risk to sight/ life (e.g. penetrating injures/globe rupture/orbital haemorrhage/burns infection-fractures/ lid lacerations)	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  b) This Prioritisation is about 'Y Whom'. Space does not allow every procedure to be listed under every	Ischaemia  Necrotising fasciitis  Bleeding (not responding to conservative Rx)  Post-operative complications (e.g. anastomotic leaks/ bleeding)  Dislocated joints  Neonate - Mixing lesions -  TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care -  Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe preeclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home/r 12-14 weeks where procedure not provided by local NHS beyond this)  Cases where cervical preparation has been administered (misoprostol/osmotic dilators/mifepristone)  Feticide (approaching legal limit)  Vitreoretinal	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine - a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	e) Safeguarding ito medical Rx.  Paediatric/adolescent  Imperforate hymen - incision and drainage  e) Safeguarding issues de tonside attending with trauma and acute surgical problems (e.g. NAI/domestic violence/	obstructive lesions - Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/ occlusion)	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)  PLEASE NOTE: More detailed specialty specific guidance can be found on the NHSE website https://www.england.nhs.u	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (requiring evacuations hysterectomy)  Acute risk to sight/ life (e.g. penetrating injures/globe rupture/orbital haemorrhage/burns infection-fractures/ lid lacerations)	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  b) This Prioritisation is about 'WHEN and not Bay Whom'. Salow devery placed under every specialty performing it and it DOES NOT indicate	Ischaemia Necrotising fasciitis Bleeding (not responding to conservative Rx) Post-operative complications (e.g. anastomotic leaks/ bleeding)  Dislocated joints  Neonate - Mixing lesions - TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care - Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe pre-eclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home)/ 12-14 weeks where procedure not provided by local NHS beyond this)  Cases where cervical preparation has been administered (misoprostol/osmotic dilators/mifepristone) Feticide (approaching legal limit)  Vitreoretinal	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine - a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	e) Safeguarding isoneside attending with trauma and acute surgical problems (e.g. NAI/	obstructive lesions - Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/ occlusion)	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)  PLEASE NOTE: More detailed specialty specific guidance can be found on the NHSE website https://www.england.nhs.uk/coronavirus/publication/	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (requiring evacuations hysterectomy)  Acute risk to sight/ life (e.g. penetrating injures/globe rupture/orbital haemorrhage/burns infection-fractures/ lid lacerations)	Fractures - Open Neurovascular compromise +/-Skin compromise  Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  Oncology - Ruthenium plaque removal.  b) This Prioritisation is about 'WHEN and not BY Whom'. Space does not allow every laparotomy/ hysterectomy  oncology - Ruthenium plaque removal.	Ischaemia Necrotising fasciitis Bleeding (not responding to conservative Rx) Post-operative complications (e.g. anastomotic leaks/ bleeding)  Dislocated joints  Neonate - Mixing lesions - TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care - Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe pre-eclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home)/ 12-14 weeks where procedure not provided by local NHS beyond this)  Cases where cervical preparation has been administered (misoprostol/osmotic dilators/mifepristone) Feticide (approaching legal limit)  Vitreoretinal	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine - a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	e) Safeguarding isometric/adolescent  Imperforate hymen - incision and drainage  e) Safeguarding issues must be considered in all those attending with trauma and acute surgical problems (e.g. NAI/domestic violence/abuse of the	obstructive lesions - Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/ occlusion)	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)  PLEASE NOTE: More detailed specialty specific guidance can be found on the NHSE website https://www.england.nhs.uk/coronavirus/publication/	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (requiring evacuations hysterectomy)  Acute risk to sight/ life (e.g. penetrating injures/globe rupture/orbital haemorrhage/burns infection-fractures/ lid lacerations)	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  Oncology - Ruthenium plaque removal.  b) This Prioritisation is about 'WHEN and not BY Whom'. Salow deer laparotomy/ hysterectomy  oncology - Ruthenium plaque removal.	Ischaemia Necrotising fasciitis Bleeding (not responding to conservative Rx) Post-operative complications (e.g. anastomotic leaks/ bleeding)  Dislocated joints  Neonate - Mixing lesions - TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care - Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe pre-eclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home)/ 12-14 weeks where procedure not provided by local NHS beyond this)  Cases where cervical preparation has been administered (misoprostol/osmotic dilators/mifepristone) Feticide (approaching legal limit)  Vitreoretinal	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine - a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	e) Safeguarding isometric/adolescent  Imperforate hymen - incision and drainage  e) Safeguarding issues must be considered in all those attending with trauma and acute surgical problems (e.g. NAI/domestic violence/abuse of the	obstructive lesions - Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/ occlusion)	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				
Obstetrics and Gynaecology (including urogynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)  PLEASE NOTE: More detailed specialty specific guidance can be found on the NHSE website https://www.england.nhs.uk/coronavirus/publication/	Septic arthritis/ osteomyelitis  Neonate - Left heart obstructive lesions - HLHS (restrictive/ intact atrial septum)  Critical aortic stenosis/coarctation (unresponsive to medical Rx)  Laparotomy/ Laparoscopy  Miscarriage with bleeding requiring surgical control and unstable  Torted/ruptured ovary/pelvic mass  Pelvic/genital tract sepsis  Bleeding  Necrotising fasciitis  Genital tract trauma - (e.g. vaginal tear/ pelvo-vaginal haematoma)  Ectopic pregnancy  Complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (complications of TOP  Molar pregnancy (requiring evacuations hysterectomy)  Acute risk to sight/ life (e.g. penetrating injures/globe rupture/orbital haemorrhage/burns infection-fractures/ lid lacerations)	Fractures - Open Neurovascular compromise +/-Skin compromise Neonate - Right heart obstructive lesions - PA-IVS PA-VSD Tetralogy of Fallot Critical pulmonary stenosis (not responding to medical Rx)  Pregnancy/Delivery Emergency Caesarean Instrumental delivery Perineal repair Manual removal of placenta Cervical cerclage Emergency laparotomy/ hysterectomy  Oncology - Ruthenium plaque removal.	Ischaemia Necrotising fasciitis Bleeding (not responding to conservative Rx) Post-operative complications (e.g. anastomotic leaks/ bleeding)  Dislocated joints  Neonate - Mixing lesions - TGA (hypoxaemia for BAS/Intact IVS for ASO)  TAPVD (clinically obstructed)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)  Early pregnancy and abortion care - Miscarriage - bleeding and unstable  Maternal compromise (e.g., sepsis, chorioamnionitis, severe pre-eclampsia, etc.)  Approaching legal threshold (23+6 weeks for all/9+6 weeks [England & Wales]/11+6 weeks [Scotland] for medical abortion at home)/ 12-14 weeks where procedure not provided by local NHS beyond this)  Cases where cervical preparation has been administered (misoprostol/osmotic dilators/mifepristone) Feticide (approaching legal limit)  Vitreoretinal	Neonate - Shunt/ stent -  Profound hypoxaemia/ occlusion/ thrombosis)  Reproductive medicine - a) Males - sperm storage before acute sterilisation b) Females - before acute sterilisation - i) Oocyte collection (n.b. must be 36hrs after the trigger) ii) Ovarian tissue storage	e) Safeguarding isometric/adolescent  Imperforate hymen - incision and drainage  e) Safeguarding issues must be considered in all those attending with trauma and acute surgical problems (e.g. NAI/domestic violence/abuse of the	obstructive lesions - Tetralogy of Fallot (cyanotic spells unresponsive to medical Rx)  Shunt/stent dependent pulmonary blood flow (pre BCPC/pre biV repair with profound hypoxaemia/ thrombosis/ occlusion)	lesions -  Aortic (haemodynamically	obstructive lesions -  MV prosthesis	Aortic (haemodynamically				

Priority 1b - Proced	-										
General surgery (including	Laparotomy -	Perianal abscess/ other infection - not	Urgent nutrition	- see notes below) Failed conservative management of	Breast sepsis - without necrosis	Upper GI endoscopy for foreign body	Acute gastric band slippage/erosion.				
oesophago-gastric, HPB, coloproctology, breast, endocrine)	obstruction - not responding to conservative Rx.  Colectomy for acute severe ulcerative colitis - not responding to conservative Rx  Bowel obstruction not suitable for	responding to conservative Rx.	Enteral nutrition access Revision Bariatric Surgery	localised intra- peritoneal infection	unresponsive to conservative Rx	removal	Acutely symptomatic internal hernia.				
OMFS	Facial fractures - not suitable for conservative Rx										
Reconstructive plastic surgery including burns and hands	Burns - requiring resuscitation.	Burns - full thickness/deep dermal requiring debridement and closure	Burns - mid/deep dermal with exposure of deep structures likely/ infection	Soft tissue infection - all sites (especially closed compartments/ joints/prostheses) not responding to conservative Rx	Delayed primary closure of open wound/fracture - any site	Primary tendon/ nerve repair - all sites.	Unstable closed fractures or joint injuries - unsuitable for conservative Rx	Secondary closure of washed out open wound/ fracture - any site	Finger tip/nail bed repair/terminalisation	Major limb trauma reconstruction unsuitable for conservative Rx	Brachial plexus/ major peripheral nerve injury - Associated with major vessel injury
Urology	Upper urinary tract obstruction	Renal stones - pain/ impairment not responsive to conservative Rx	Penile fracture	Infected prosthesis - penile/testicular/ ureteric stent							
T & O (including spinal surgery)	Tibial fracture - high energy/displaced, unstable shaft/	Fractures - pathological and peri-prosthetic	Unstable articular fractures that will result in severe disability without operative fixation	Non-hip lower limb frailty fractures requiring fixation to mobilise patient	Spinal Trauma requiring stabilisation without neurological involvement						
ENT	Other foreign body in nose	Orbital decompression	Acute mastoiditis and other middle ear conditions not responding to conservative Rx	Traumatic/ cholesteatoma related facial nerve palsy	Traumatic injury to the pinna	Lymph node biopsy - lymphoma where core biopsy inadequate.	Head and neck sepsis - not responding to conservative Rx.	MDT directed cancer debulking/biopsy - Microlaryngoscopy +/- laser	Vocal Cord medialisation for severe aspiration	Compound/complex fractures of the nose and sinuses	Insertion for patients with meningitis.  Removal of infected implant not responding to
Neurosurgery (including spinal surgery)	Depressed skull fracture	Traumatic brain injury - not responding to conservative Rx - neurological compromise	Intracranial haemorrhage - no longer responding to conservative Rx	Acute raised Intra cranial pressure/ hydrocephalus (recoverable stroke/ tumour) - no longer responding to conservative Rx	Battery change for spinal/deep brain/ epilepsy stimulators/pumps	MDT directed paediatric brain tumour surgery					conservative Rx.
Cardiothoracic surgery	Empyema not responding to Rx	Coronary Artery Disease - Unstable/ Rest ECG changes and not reposing to conservative Rx	Aortic Valve Disease - Deteriorating Symptoms / Haemodynamically unstable	Mitral Valve Disease - Deteriorating Symptoms / Haemodynamically unstable	Myxoma - Emboli/ Haemodynamically unstable	Chest Trauma					
Vascular surgery	Acute on chronic limb ischaemia	Symptomatic carotid disease	Amputation for limb ischaemia	Diabetic foot sepsis - not responding to conservative Rx	Blocked AV fistula						
Paediatric general and urological surgery (see also urology)	Neonatal Malformations -  Duodenal Atresia,  Small bowel obstruction  Large bowel obstruction  Congenital Diaphragmatic Hernia  Congenital Pulmonary Airway Malformations (CPAMS) - respiratory compromise	Laparotomy - small bowel obstruction not responding to conservative Rx	Laparotomy - Colectomy for colitis (Ulcerative Colitis/ Hirschsprung's) not responding to conservative Rx	Soft tissue infection - any site not responding to conservative Rx	Central Venous Line insertion for Oncology/Enteral nutrition/Access for antibiotics/Dialysis	Malignant tumour/ Lymph node biopsy	Pyloromyotomy	Peritoneal Dialysis Catheter Insertion	Resection of Posterior Urethral Valves	Exstrophy - Primary bladder closure	Hydronephrosis - Rapid progression
Paediatric orthopaedic surgery (including spinal surgery)	Slipped Upper Femoral Epiphysis	Fractures -  Displaced articular/ peri-articular  Forearm  Femoral	Exposed metalwork								
Paediatric cardiac surgery Obstetrics and	Laparotomy/	Incision + drainage/	Miscarriage -	Abortion -	Fistula repair -	MDT Directed EUA	Hysteroscopy -				
Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)	Laparoscopy  Pelvic collection/ tubo-ovarian abscess (not responding to conservative treatment, incl. interventional radiology)  Ectopic pregnancy (stable patient)  Evacuation of haematoma/Repair wound dehiscence/ Evisceration/ Incisional hernia  Pelvic pain >48 hours  Bowel obstruction - Cancer not responding to conservative Rx.	marsupialisation - Bartholin's abscess	Patient stable - case selection	All cases -NOS  (From NICE 2019: ensure minimum delay and provide within 1 week)	Recto-vaginal/ Bladder-vagina	and insertion of fiducial markers - Cervical cancer staging and planning	PMB with thickened endometrium + not amenable to outpatient sampling				
Ophthalmology		Vitreoretinal -  Laser/cryotherapy - retinal tear  Vitrectomy - i) dropped lens nucleus after cataract surgery  ii) Detachment - macular on/recently off	Cornea - Corneal transplant/ glueing Amniotic membrane graft - threat to sight		Glaucoma -  Acute - i) Laser PI ii) Unresponsive to medical Rx/laser  Secondary - Drainage/diode laser - imminent risk to sight	Paediatrics -  Retinopathy of prematurity - retinal -laser/intravitreal injection  Examination under anaesthesia - potential threat to sight  (see also trauma)	Medical -  Retina - Periocular/intravitreal steroids for inflammatory eye disease  Temporal artery biopsy				
PLEASE NOTE: More detailed specialty specific guidance can be found on the NHSE website https:// www.england.nhs.u k/coronavirus/ publication/ specialty-guides/	a) THIS DOCUMENT WILL BE REVIEWED MONTHLY	b) This Prioritisation is about 'WHEN and not BY Whom'. Space does not allow every procedure to be listed under every specialty performing it and it DOES NOT indicate primacy of ANY specialty legitimately performing any procedure within their listed competencies.		Surgery in Paediatric patients is included in the	e) This RPM matrix is to be used ONLY to assess patients in the SAME priority band.  **Priority band**  **Pr	f) Patients in p1b MUST be regularly reviewed clinically and re-prioritised to; p1a if their clinical condition deteriorates. p2 If their clinical condition improves and stabilises.					

Priority 2 - procedu	res to be performed i	n < 1 months.												
(n.b. This prioritisating General surgery (including oesophago-gastric, HPB, coloproctology, breast, endocrine)	stricture/fistula not responsive to endoscopic/medical Rx	MDT Directed hepatobiliary/ pancreatic/	mg the Covid19 Crisis  MDT Directed resection/diversion for highly symptomatic cancer.	MDT Directed thyroid/parathyroid cancer surgery	Thyrotoxicosis - Not responding to conservative Rx. (including orbital surgery for impending sight loss)	Parathyroidectomy - calcium >3.0mmol/l and/or not responding to conservative Rx, especially pregnancy/post-transplant/repeated admission.	MDT Directed adrenal cancer surgery	Adrenalectomy - pathology not responding to medical Rx (e.g. Cushing's/ phaeochromocytom a)	MDT Directed breast cancer surgery and IBR, if appropriate according to local fitness criteria -  ER negative  Her2+  Pre-menopausal  ER+ with higher risk (i.e., Grade 3, Low		MDT directed bariatric surgery  As part of cancer/ transplant treatment.			
OMFS	oropharyngeal/tonsil/	Facial Fractures causing diplopia/ occlusal problems	Mandibular/maxillary orthognathic surgery - airway compromise unresponsive to conservative Rx AND unsuitable for tracheostomy - adults and children	Adult and paediatric	Craniofacial - ocular complication/Raised Intracranial Pressure				ER, node +ve)					
Reconstructive plastic surgery including burns and hands		Removal of prosthesis - unresponsive to conservative Rx.	Burns - Reconstruction  Eyelid closure problems  Severe microstomia  Joint and neck contracture		Skin cancer -  Primary resection directed by appropriate skin cancer specialist	Skin cancer -  MDT Directed further resection  Re-excision according to national guidelines  SLNB and all completion lymphadenectomies  Electrochemotherapy	Closed injury - not suitable for observation.  Exploration for paresis/pain/sensory impairment	Eyelid closure/	Neonate accessory digit excision (narrow pedicle/ vascular compromise/ infection/pain)					
Urology		MDT directed penile cancer surgery including inguinal node surgery.	MDT directed bladder cancer surgery - invading bladder muscle.	cancer surgery - not		ILP MDT directed bladder Cancer surgery - high risk carcinoma-in- situ.		Acute Urinary Retention - Bladder neck stenosis post RARP.	Partial Nephrectomy - single kidney	Visible haematuria - investigation				
T & O (including spinal surgery)		Solitary metastasis surgery - any site.	MDT Directed destructive bone lesion surgery with risk of fracture (e.g. Giant cell tumour)	Displaced, intra-	Knee extensor disruption (including fractured, displaced patella)	Tendon rupture - any sites	Locked joints - any site	Peripheral Nerve Decompression - any site (pain/ weakness/muscle wasting - not responding to conservative Rx)	Arthroplasty - any site where delay will prejudice outcome	Spinal surgery - degenerative conditions with progressive neurology/ neurological deficit				
ENT	EUA/biopsy for malignancy - hypopharynx/larynx	MDT directed nasopharyngeal/ laryngeal surgery for malignancy	MDT directed oropharyngeal r surgery for malignancy		Baro-trauma perilymph fistula	Organic foreign bodies in the ear.	MDT directed treatment of small, high grade salivary cancers.	MDT directed treatment of sinus cancers threatening sight	Treatment of pharyngeal/oesophageal/airway stricture	Mucocoele with recurrent infection/ visual disturbance		Reduction of nasal fracture (NOS) - ideally inside 14 days according to local capacity	Cochlear implant -  Pre -lingually deafened children  Post- meningitis  Device failure - no hearing	Airway compromise - NOS
Neurosurgery (including spinal surgery)	MDT directed brain tumour surgery (including for metastases)	MDT directed spinal tumour surgery	Acute/chronic pain syndromes - (e.g. trigeminal neuralgia) - unresponsive to conservative Rx											
Cardiothoracic surgery	MDT directed treatment of resectable Non- Small Cell Lung Cancer	Unstable Non ST elevated MI	Aortic stenosis	Unstable coronary	Any deteriorating heart condition	Pneumothorax not responding to conservative Rx								
Vascular surgery	Chronic severe limb ischaemia - no neurology	AAA >7cms diameter												
Paediatric general and urological surgery (see also urology)	Laparotomy or Stoma Closure to manage intestinal failure with liver disease / complications	Infant with Billary Atresia - bladder exstrophy	Inguinal hernia under 3/12 of age	surgery for Nephroblastoma/	Crohn's Disease - stricture/fistula/ optimise medication/ nutrition	Gastrostomy for nutritional support.	Fundoplication for GOR for previous life threatening complication/ repeated aspirations	Renal Stent Removal/Exchange	Vesico-ureteric reflux - case selection	Circumcision for severe BXO/meatal pathology	Recurrent UTIs - renal scarring/ hypertension	Non-functioning renal tract with infection – not responding to conservative Rx		Renal Calculi - recurrent symptoms/ renal impairment
Paediatric orthopaedic surgery	•	MDT Directed Suspected, aggressive, benign bone tumour	Meniscal repair											
Paediatric cardiac surgery	obstructive lesions - Aortic stenosis (valvuloplasty/ valvotomy)  Coarctation (case selection of approach and timing)  HLHS (Norwood/ Hybrid)	heart obstructive lesions - PA-IVS (case selection RF perforation/ductal stent/shunt)	Neonate - Mixing lesions -  TGA (Intact IVS for ASO/VSD with mixing for ASO+VSD)  TAPVD (echo evidence of obstruction)  Common arterial trunk (excess pulmonary blood flow, truncal regurgitation not responding to medical Rx)		Neonate - ALCAPA - (Optimise medical Rx)	Infant - Left heart obstructive lesions -  LVOTO (impaired function/symptoms)  Aortic stenosis - (impaired function/symptoms)  Coarctation (Impaired function)	obstructive lesions - Tetralogy of Fallot (Cyanotic spells cyanosis <80%) Shunt/stent dependent pulmonary blood flow (pre BCPC -increasing cyanosis/ shunt/stent stenosis	VSD (FTT, not responding to medical Rx, >6 months of age)  AVSD (FTT, not responding to medical Rx, assessment of	function)	Child - Left heart obstructive lesions - LVOTO (impaired function/symptoms)  Aortic stenosis (impaired function/symptoms)  MV prosthesis (increased gradient/raised RVP)	obstructive lesions -  RV-RA conduit (impaired function/ >systemic RVP)	Child - Regurgitant lesions -  Mitral (not responding to medical Rx/raised RVP)  Aortic (impaired function/symptoms)	Child - Fortan candidate - (increasing cyanosis/ symptoms) prioritise >5yrs years old	
Obstetrics and Gynaecology (including urogynaecology, pregnancy, delivery, and reproductive medicine)	MDT redirected cancer surgery –  Staging  Vulval/vaginal (incl. WLE for high grade Vulval Intraepithelial Neoplasia (VIN), Vaginal Intraepithelial Neoplasia)  Uterine/Ovarian/Cervical (incl. Intraepithelial Neoplasia/early stage cancer)  Recurrent Gynaecological cancer	Hysteroscopy +/- endometrial Bx for endometrial hyperplasia/cancer	Paediatric and adolescent –  Non-obstructive vaginal septum/ septate hymen  EUA/vaginoscopy for suspected vaginal abnormality											
	Ocular/ocular surface tumours  Enucleation - advanced melanoma/other malignancies/ tumours  Intravitreal injections - radiation maculopathy +/-	Vitreoretinal - Re-do retinal detachment Vitrectomy - i) Retinal detachment - macular off > 2 weeks ii) Acute vitreous haemorrhage - suspected retinal break/unknown iii) Dislocated lens implant with poor vision in other eye	Adnexal -  Protect ocular surface  MDT directed treatment for eyelid orbital tumours	sight (also see cataract)  i) Very high IOP	Cataract - Intumescent extraction  Angle closure glaucoma - threat to sight	Cornea -  Amniotic membrane graft - non-healing ulcer (see also paediatrics)	Paediatrics - Congenital cataract Keratoplasty for congenital corneal opacity Superficial keratectomy - atopic plaque Drainage surgery - glaucoma (< 2 weeks) Surgery/plaque -retinoblastoma Brow suspension - risk of developing amblyopia	Medical retina –  Intravitreal injections for wet, age related macular degeneration >2/52  Laser for active/ progressive neovascularisation  Periocular and intravitreal steroid injection for macular oedema	Strabismus – Sudden loss of binocularity					
PLEASE NOTE: More detailed specialty specific guidance can be found on the NHSE website https:// www.england.nhs.u k/coronavirus/ publication/ specialty-guides/	DOCUMENT WILL BE REVIEWED MONTHLY	b) This Prioritisation is about 'WHEN and not BY Whom'. Space does not allow every procedure to be listed under every specialty performing it and it DOES NOT indicate primacy of ANY specialty legitimately performing any procedure within their listed competencies.		Surgery in Paediatric patients is included in the guidance above.	e) Safeguarding issues must be considered in all those attending with trauma and acute surgical problems (e.g. NAI/domestic violence/abuse of the vulnerable)		Meanur Principles (Inc.) 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1	who have not been						

General surgery including besophago-gastric, HPB, coloproctology, breast, endocrine)	MDT directed cancer surgery - Colon cancer/liver metastases	-	MDT Directed thyroid cancer surgery - including diagnostic	- see notes below)  Colectomy/ proctectomy for colitis refractory to medical Rx		cancer surgery and IBR, if appropriate		Cholecystectomy - post acute pancreatitis	Hernia - presenting with complications that have previously		MDT directed full thickness rectal prolapse surgery	MDT Directed bariatric surgery
Desophago-gastric, HPB, coloproctology, preast, endocrine)	Colon cancer/liver		thyroid cancer surgery - including	proctectomy for colitis refractory to	fistulae (incl. perianal	IBR, if appropriate		post acute	with complications	- symptomatic renal		bariatric surgery
coloproctology, preast, endocrine)		ператоріliary	uaanostic	medical Ry		Locaria :	I		COLLIN III	MO T IT I' '	promped dangery	Ciamin's art are
		Pancreatic	lobectomy.	(excluding acute, severe colitis treated	Crohn's)	according to local fitness criteria.	>4cm<6cm) with		settled with conservative Rx	responding to conservative Rx.		Significant/multiple end organ failure.
	Rectal cancer/liver metastases	Oesophagogastric		urgently)		Pre-menopausal ER+ (Grade 1-2)	(Cortisol/androgen)					To facilitate MSK/ Hernia Surgery liste
	Multi-visceral resections for locally advanced colon cancer	Neuroendocrine tumour				Post-menopausal ER+ with higher risk (i.e., Grade 3, low ER or node +ve)	metastases - progressing on scan at 3/12.					in p3  Overdue balloon removal.
	Salvage surgery for recurrent anal cancer					High grade DCIS  Risk reducing						Revision to stop excessive weight loss/comorbidities.
OMFS I		MDT directed salivary gland tumours (low grade).	Cleft lip - Alveolar bone grafting (Prior to canine eruption)			surgery in gene carriers.						
,	well differentiated with no metastases.											
olastic surgery ncluding burns	Burns - Reconstruction  Microstomia	Limb contractures	Secondary cleft and non-cleft speech surgery - to avoid breaching 5 yrs of age	Primary cleft palate Repair - to avoid breaching 13 months of age	Brachial plexus/ major peripheral nerve injury - MDT Directed	Facial Palsy -  Dense facial palsy inside 12/12 from injury	Congenital hand anomaly where delay will compromise outcome.	MDT directed surgery for major upper limb functional impairment				
	Joint contracture Neck contracture		ago		i) re-animation +/- joint stabilisation  ii) Exploration for life altering pain not responding to conservative Rx.  iii) Revision surgery for major functional							
		Stent removal/	Haematuria -	MDT directed	impairment.  MDT Directed penile							
;	prostate cancer surgery - high/ intermediate risk	exchange	investigation for non-visible (including paediatric)	bladder cancer surgery (not invading muscle)	cancer surgery (low grade and premalignant).							
	Hip Avascular Necrosis (night pain/	Frozen shoulder - severe and not	Tendon reconstruction/	Revision surgery	MDT Directed Benign bone/soft	MDT Directed primary sarcoma	Arthroscopic removal of joint	other reconstruction	Removal of metalwork	Spinal Surgery – Injection/		
	collapse of the joint/	responding to conservative Rx	tenodesis - any site	Loosening without impending fracture. Recurrent joint instability	tissue lesion excision biopsy - not otherwise specified	plus metastases	loose body (Reversible symptoms preventing work)	555454011	e.g. across joints.	decompressive surgery for intractable radiculopathy.		
ENT (	CSF fistula repair	Expanding	Cochlear implant in			· ·	Sinus surgery with					
		mucocoele without infection/NOS	pre-verbal profound hearing loss where delay will impact on long term outcome.	otological cancer surgery.	and papilloma resection (laser/ microdebrider/ coblation/steel)	treatment of pharyngeal pouch with severe dysphagia	complication of infection					
Neurosurgery												
including spinal surgery)	Otels N 5											
surgery  /ascular surgery	Stable Non ST Elevation MI AAA >5.5cm and <7cm diameter											
and urological surgery (see also	Congenital Malformations with delayed Management -	Inguinal hernia 3-12 mths of age	Gastrostomy for Failure To Thrive (FTT)	Interval appendicectomy for recurrent symptoms	Cholesystectomy	Fundoplication for GOR - failure to thrive	Orchidopexy for undescended testis	Daytime urinary incontinence - obstructive cause suspected.	Penile anomalies - (e.g., mega prepuce but <b>not</b> hypospadias.)	Varicocoele/ Hydrocoele - large + symptomatic.		
	Hirschsprung's Disease initially managed with washouts.											
orthopaedic surgery (including	Developmental Dislocation of the Hip (DDH) - Primary joint stabilisation	Congenital Talipes Equino Varus (CTEV) - Initial management including tenotmies	Limb length discrepancy/ malalignment	Childhood/ adolescent spinal deformity								
Paediatric cardiac												
Surgery Obstetrics and Gynaecology	Urogynaecology -	MDT Directed cancer treatment -	Hysteroscopic/ Laparoscopic/Open	BSO/salpingectomy	Hysterectomy - risk reducing for Lynch	Fertility -	Paediatric and adolescent – MDT					
including urogynaecology,	Suprapubic catheter change	Cervical (Stage Ia1)	Myomectomy/ Hysterectomy/	Risk reducing for	Syndrome	Pelvic pathology effecting fertility	directed					
oregnancy, delivery, and	Prolapse - bleeding/	at 6-8/52 pending MDT outcome	Endometrial ablation			(e.g., Fibroids/ Hydrosalpinx/ Endometriosis/	Laparoscopic excision of obstructed uterine					
medicine)	ulceration/proci dentia/vault inversion	Repeat conisation - Any age/High grade	conservative Rx)	Complex ovarian		Uterine septum/ Adhesions)	horn					
	Genitourinary fistula	pre-cancer with pt. >50 yrs of age)	Fibroids/Heavy menstrual bleeding (significant anaemia	cyst - low risk of malignancy		Couples/individuals where the woman	Vaginal reconstruction for agenesis with					
		Simple hysterectomy following local	1, 0			has low ovarian reserve >40 years	menstrual obstruction					
		conisation (LLETZ)  Low volume cancer	Endometriosis - a) Severe symptoms			old.						
		completely excised at loop excision.	unresponsive to medical Rx									
		Low grade uterine cancer managed conservatively with LNG-IUS and/or oral progestogens.	b) Bowel/ureteric obstruction - failed/ unsuitable for stenting)									
Ophthalmology	Vitreoretinal -	Adnexal -	Glaucoma -	Cataract -	Cornea -	Paediatrics -	Medical –	Strabismus –				
	Some Macular holes	_	Drainage - not otherwise specified	Surgery/YAG laser i) Binocular vision	Cross-linking - rapidly progressive/	Retinal laser/ cryotherapy/	Diabetic macula/ retinal vein/branch vein occlusion	Development binocularity in				
1	Vitrectomy - i) Vitreous haemorrhage/	Entropion/Ectropion - ocular surface damage	Selected laser trabeculoplasty	<6/60/severely disabled (e.g. cannot work)	very thin cornea	intravitreal injections - Retinal vascular conditions	i.Intravitreal injections ii.Macular laser	infantile squint Surgery or botulinum				
1	tractional retinal detachment	Eye removal - Non-	ι. αροσαίσριασιγ	ii) limiting		Capsulotomy - visual	ii.Macular laser Photodynamic	injection for severe diplopia (e.g. cannot				
	ii) silicone oil removal -	malignant/low threat to health		management of other conditions - threat to sight		axis opacity following congenital cataract surgery	laser for central serous	work)				
	complications	Botulism injections for disabling		<del></del>		Removal of loose	chorioretinopathy					
		blepharospasm				corneal sutures in children (see also strabismus)						
		b) This	c) Any delay in	d) Other Specialist	e) Safeguarding	f) Private sector	g) This RPM matrix					
specialty specific guidance can be	BE REVIEWED	Prioritisation is about 'WHEN and not BY Whom'.	,	Surgery in Paediatric patients is included in the	those attending	procedures should be considered on	in the SAME	treated MUST be reviewed clinically				
ound on the NHSE website https://www.england.nhs.u		Space does not allow every procedure to be	and life threatening conditions, may lead to adverse	guidance above.	with trauma and acute surgical problems (e.g. NAI/	merit and on a case-by-case basis.	priority band.	at most 3/12 from being listed and re- prioritised as				
c/coronavirus/ oublication/ specialty-guides/		listed under every specialty performing it and it	outcomes.		domestic violence/ abuse of the vulnerable)	Procedures with a known functional	The second secon	necessary.				
		DOES NOT indicate primacy of ANY specialty				benefit should be prioritised where possible.	Contract pursue, Contra					
		legitimately				A detailed risk	Comment Management Comment Com					
		performing any					sames that dis real.  Self-like an elization law suppressurate  Law like an elization law suppressurate  same places law like and  same places law l					
		performing any procedure within their listed competencies.				analysis should be undertaken and consideration given to any potential	Martin Andrews (Martin Andrews					

	_	res to be performed		ing the Covid19 Crisis	- see notes below											
General surgery including	MDT directed breast cancer surgery			Diverting ileostomy closure	Uncomplicated incisional hernias	Reversal of Hartmann's	Non-urgent proctology	Transanal/rectal resection of benign	Benign breast disease	Cholecystectomy - after biliary colic/	Other benign upper UGI conditions (e.g.	Oesophagogastric	Other benign thyroid/parathyroid	Other adrenal disease -	Abdominal wall reconstruction	MDT Directed bariatric surgery
nesdang besophago-gastric, IPB, coloproctology, breast, endocrine)		Benign colonic polyp Completion proctocolectomy for IBD	0	Cioduic		procedure	procedures	rectal polyps.	Delayed breast reconstruction, if appropriate according to local fitness criteria.  Revision of breast reconstruction,	cholecystitis.	gallstones/other Benign disease).	Tellux Surgery	disease - uncomplicated	uncomplicated	reconstruction	Lesser degrees end organ failur
	DCIS (intermediate and low risk)															
OMFS	All orthognathic Surgery	Dental extractions - adult and paediatric	1	Facial deformity - Post-traumatic/ Cancer treatment	Benign dental lesions - mandible/ maxilla	Temporo-mandibular joint surgery										
Reconstructive plastic surgery including burns and hands	Burns - other contractures/scars	Limb trauma sequelae/scarring - other reconstruction	Breast reconstruction - NOS	All cleft lip and palate surgery - NOS	Basal Cell Carcinoma - any site not compromising vital structures	Excision of benign lesions	NHS Cosmetic Surgery									
Urology		Andrology/GU Surgery  Erectile dysfunction  Male fertility surgery  Urethral stricture  Gender reassignment.	Endourology -  Uncomplicated stones/  Percutaneous nephrolithotomy/  Pelviureteric obstruction	MDT directed prostate cancer surgery (low risk)	MDT directed bladder cancer surgery - superficial transitional cell cancer	Uncomplicated small/intermediate renal lesions	Uncomplicated small/intermediate testicular lesions	Bladder outflow surgery	Benign penoscrotal surgery	Renal stones - asymptomatic	Uncomplicated small/intermediate renal lesions					
T & O (including spinal surgery)	Arthroplasty/ arthrodesis - not otherwise specified,	Hand and Upper limb surgery - Not otherwise specified	Metalwork removal	Degenerative spinal disease - no neurological compromise/ refractory pain	Adult spinal deformity surgery with progression											
ENT	All other Rhinology (septoplasty/ septorhinoplasty/ turbinate surgery/ sinus surgery [NOS])	Cholesteatoma - uncomplicated.	Chronic suppurative otitis media	All Ossicular Surgery/Middle ear implants	Tympanopasty	Grommets	Meatoplasty	Vestibular Surgery	Non-organic foreign body (except button batteries)	Cochlear Implants - other	Uncomplicated nasal fracture	Micro Laryngoscopy - benign vocal fold/ cord conditions (e.g. polyp/cyst/ ectasia/paralysis)		Routine procedures for pharyngeal pouch		
Neurosurgery (including spinal surgery)	Congenital spinal deformity - no neurological compromise/ refractory pain	Movement disorder implants	Lesioning/epilepsy surgery	Normal pressure hydrocephalus	Slow growing brain tumours - no neurological compromise	Slow growing spinal tumours - no neurological compromise	Benign intracranial arteriovenous malformations/ tumours) - no neurological compromise	Paediatric craniofacial surgery not compromising vision/neurology/ raised ICP	-							
Cardiothoracic surgery Vascular surgery	Stable coronary disease  Vein surgery	Thoracic outlet	Claudication													
Paediatric general and urological surgery (see also urology)	Anoplasty/Posterior Sagittal Ano- Rectoplasty (PSARP) - after obstruction relieved	mths of age)	haemoglobinopathy	Cholecystectomy - after biliary colic/ cholecystitis	Stoma Closure	Benign lesion excision	Hypospadias repair (around 12 mths of age)	Cosmetic foreskin issues/ritual circumcision.	Epispadias	Gender dysphoria	Daytime urinary incontinence - all children >7yrs	Asymptomatic hyrocoele				
Paediatric orthopaedic surgery (including spinal surgery)	Developmental Dislocation of the Hip (DDH) - Secondary joint reconstruction	Congenital Talipes Equino Varus (CTEV) - Late presenting/relapsed	Spasticity management	Corrective surgery for established deformity	Reconstruction for established joint instability (e.g. ACL/Lateral ligament)	Metalwork removal										
Paediatric cardiac	reconstruction															
surgery Obstetrics and Gynaecology	Urogynaecology	Hysteroscopy -	Laparoscopy -	Laparoscopic/Open myomectomy -	Laparoscopic/Open or Vaginal	Laparoscopic/Open cystectomy/	Symptomatic lower genital tract lesions	Closure of Stoma	Fertility -	Paediatric and adolescent –						
(including urogynaecology, pregnancy, delivery, and reproductive medicine)	Incontinence surgery Prolapse surgery	bleeding/ Reproductive failure (e.g. Levonorgestero	Tubal factor infertility +/- symptomatic tubal disease	Fibroids not causing anaemia	hysterectomy - Abnormal uterine bleeding Pain Symptomatic fibroids +/- endometrial hyperplasia	Oophorectomy -  Ovarian cysts > 5 cm with a benign RMI)	(e.g. uninfected Bartholin's cyst)		Pelvic pathology effecting fertility (e.g., Fibroids/ Hydrosalpinx/ Endometriosis/ Uterine septum/ Adhesions)  Couples/individuals where the woman has normal ovarian reserve <40 years old	Vaginal reconstruction (NOS)  Clitoral reduction for differences in sex						
Ophthalmology	Vitrectomy - i) Macular epiretinal membrane ii) Silicone oil removal - not otherwise specified (NOS) iii) Other surgery - NOS	Adnexal -  Dacrocystorhinosto my  Other surgery - NOS	Cataract - YAG laser capsulotomy Significant binocular visual reduction Other surgery - NOS	Cross-linking - progressive keratoconus  Corneal graft - significant binocular			pre-proliferative diabetic retinopathy	Glaucoma – PI laser for narrow angles								
PLEASE NOTE: More detailed specialty specific guidance can be found on the NHSE website https:// www.england.nhs.u k/coronavirus/ publication/ specialty-guides/	1	b) This Prioritisation is about 'WHEN and not BY Whom'. Space does not allow every procedure to be listed under every specialty performing it and it DOES NOT indicate primacy of ANY specialty legitimately performing any procedure within their listed competencies.		Surgery in Paediatric patients is included in the	e) Safeguarding issues must be considered in all those attending with trauma and acute surgical problems (e.g. NAI/domestic violence/abuse of the vulnerable)	aesthetic surgery procedures should be considered on merit and on a case-by-case basis.	in the SAME priority band.  Service of the priority band.  Service of the priority band of th									